arXiv:1408.4828v3 [math.DS] 7 Dec 2014

DELONE PROPERTY OF THE HOLONOMY VECTORS OF TRANSLATION SURFACES

CHENXI WU

Abstract. We answer a question by Barak Weiss on the uniform discreteness of the set of the holonomy vectors of translation surfaces.

For a translation surface M, let $S_M \subset \mathbb{R}^2$ be the set of holonomy vectors of all saddle connections of M. The following is a plot of S_M where M is a lattice surface in $H(2)$ with discriminant 13.



A way of capturing the concept of uniformity of a subset of a metric space is the concept of Deloné set.

Definition 1. [Sen06] Subset A of a metric space X is a Deloné set if:

(1) A is relatively dense, i.e. there is $R > 0$ such that any ball of radius R in X contains at least one point in A.

(2) A is uniformly discrete, i.e. there is $r > 0$ such that for any two distinct points $x, y \in A$, $d(x, y) > r$.

The Deloné property implies a quadratic upper and lower bound on the growth rate of S_M (the number of points within radius R of the origin). It is known that the growth rate of S_M does satisfy such upper and lower bounds by [Mas88] and [Mas90]. For some translation surfaces the asymptotic upper and lower bounds agree. Veech [Vee89], Eskin and Masur [EM01] showed that this is the case for Veech surfaces and generic translation surfaces respectively. Also, whether or not M is a lattice surface is determined by properties of S_M as shown in Smillie and Weiss [SW10]. Furthermore, additional properties that S_M has to satisfy are contained in the works of Athreya and Chaika [AC12, ACL13] on the distribution of angles between successive saddle connections of bounded length.

Barak Weiss asks for which translation surfaces M, is S_M a Deloné set. Here, we will show that:

Theorem 1. *If M is a lattice surface then S_M is never a Deloné set. On the other hand, there exists a non-lattice translation surface M for which S_M is a Deloné set.*

We will show that if M is a non-arithmetic lattice surface, then S_M cannot be uniformly discrete. We will also show that if M is square-tiled then S_M cannot be relatively dense. Combining these two results, we can conclude that when M is a lattice surface, S_M cannot be a Deloné set.

Theorem 2. *If M is a non-arithmetic lattice surface, then S_M is not uniformly discrete.*

Proof. Let M is a non-arithmetic lattice surface and let $r > 0$ be given. By [Vee89] we can choose a periodic direction γ of M such that in this direction M is decomposed into cylinders $C_1, \ldots, C_n$ in the direction of γ, and the width of all these cylinders are no larger than $r/4$. Because M is a lattice surface, the holonomy field [KS00] is generated by the ratios of the circumferences. Because M is not square-tiled, the holonomy field can not be $\mathbb{Q}$. Hence, there exist two numbers i and j such that the quotient of the circumferences of C_i and C_j is not in $\mathbb{Q}$. Denote the holonomy vectors of periodic geodesics corresponding to C_i and C_j by l and l'. Let h and h' be the holonomy vectors of two saddle connections α_0 and β_0 crossing C_i and C_j respectively. Let α_n be the images of α_0 under n-Dehn twists in cylinders C_i, β_n be the image of β_0 under n-Dehn twists in cylinder C_j. Given $n \in \mathbb{Z}$, both α_n and β_n are still saddle connections of M. Thus, for any integer n, the vectors $h + nl$ and $h' + nl'$ are in S_M.



the cylinder C_i



the holonomy vector of α_0 and its images under Dehn twists



the cylinder C_j



the holonomy vectors of β_0 and its images under Dehn twists

Let us write h as $h = h_1 + h_2$ and h' as $h' = h'_1 + h'_2$, where h_1, h'_1 are vectors in direction γ, and h_2, h'_2 are vectors in direction $\gamma^\perp$. Because the width of C_i and C_j are no larger than $r/4$ by assumption, $||h_2 - h'_2|| < r/2$. Because h_1, h'_1, l and l' are vectors pointing in the same direction, we can write $h_1 = al$, $h'_1 = bl$, $l' = \lambda l$. Because the quotient λ of lengths of l' and l is irrational, the set $\{m + m'\lambda : m, m' \in \mathbb{Z}\}$ is dense in $\mathbb{R}$, so there exists a pair of integers n_0 and n'_0 such that $|a - b + n_0 - n'_0\lambda| < \frac{r}{2||l||}$. Thus $||(h_1 + n_0 l) - (h'_1 + n'_0 l')|| < r/2$, and $||(h + n_0 l) - (h' + n'_0 l')|| < r/2 + r/2 = r$. We conclude that S_M contains two points for which the distance between them is less than any $r > 0$, thus S_M is not uniformly discrete. □

The above argument also works on those completely periodic surfaces such that in any given periodic direction, there are at least two closed geodesics whose length are not related by a rational multiple. Barak Weiss pointed out that the above argument implies that any

orbit in $\mathbb{R}^2$ under a lattice in $SL(2,\mathbb{R})$ is either contained in a lattice in $\mathbb{R}^2$ or not uniformly discrete.

Now we deal with the square-tiled case. This case is closely related to the classical case of the torus discussed in [HS71]. In this case, $S_M = \{(p,q) : \gcd(p,q) = 1\}$

Lemma 3. *For any positive integer N, the set $\{(p,q) \in \mathbb{Z}^2 : \gcd(p,q) \leq N\}$ is not relatively dense in $\mathbb{R}^2$.*

Proof. ([HS71]) Given $R > 0$, choose an integer $n > 2R$, and n^2 distinct prime numbers $p_{i,j}$, $1 < i,j < n$ larger than N. Let $q_i = \prod_j p_{i,j}$ and $q'_j = \prod_i p_{i,j}$. By the Chinese remainder theorem there is an integer x such that for all i, $x \equiv -i \mod q_i$, and an integer y such that for all j, $y \equiv -j \mod q'_j$. Hence for any two positive integers $i,j \leq n$, $(x+i, y+j) \in \{(p,q) \in \mathbb{Z}^2 : \gcd(p,q) > N\}$, in particular, there is a ball in $\mathbb{R}^2$ of radius R that does not contain points in the set $\{(p,q) \in \mathbb{Z}^2 : \gcd(p,q) \leq N\}$. □

Now we use the above lemma to show that if the surface M is square-tiled, S_M cannot be relatively dense.

Theorem 4. *If M is a square-tiled lattice surface then S_M is not relatively dense in $\mathbb{R}^2$.*

Proof. If M is square-tiled, we can assume that M is tiled be 1×1 squares. Let N be the number of squares that tiled M, then M is an n-fold branched cover of $T = \mathbb{R}^2/\mathbb{Z}^2$ branched at $(0,0)$. Therefore, the holonomy of any saddle connection is in $\mathbb{Z} \times \mathbb{Z}$. For any pair of coprime integers (p,q), let γ be the closed geodesic in T starting at $(0,0)$ whose holonomy is (p,q). The length of γ is $\sqrt{p^2+q^2}$. The preimage of γ in M is a graph Γ. The vertices of Γ are the preimages of $(0,0)$, while edges are the preimages of γ. The sum of the lengths of the edges of Γ is $N\sqrt{p^2+q^2}$. Any saddle connection of M in (p,q)-direction is a path on Γ without self intersection, hence The length of such a saddle connection can not be greater than $N\sqrt{p^2+q^2}$. Hence, the holonomy of such a saddle connection is of the form (sp, sq), $s \in \mathbb{Z}$ with $|s| \leq N$. Thus, $S_M \subset \{(p,q) \in \mathbb{Z}^2 : \gcd(p,q) \leq N\}$, so by Lemma 3 S_M is not relatively dense in $\mathbb{R}^2$. □

Finally, when M is not a lattice surface, S_M *can* be a Deloné set, which we will show in Example 1 below. This construction finishes the proof of Theorem 1.

Example. Let M_1 be the branched double cover of $\mathbb{R}^2/\mathbb{Z}^2$ branched at points $(0,0)$ and $(\sqrt{2}-1, \sqrt{3}-1)$. Let $\tilde{M}$ be a $\mathbb{Z}^2$-cover which is a branched double cover of $\mathbb{R}^2$ branched at $U = \mathbb{Z}^2$ and at $V = \mathbb{Z}^2 + (\sqrt{2}-1, \sqrt{3}-1)$, where the deck group action is by translation. Then saddle connections on M_1 lift to saddle connections on $\tilde{M}$, and any two lifts have the same holonomy, hence S_{M_1} is the same as $S_{\tilde{M}}$, which is the set of holonomies of line segments linking two points in $W = U \cup V$ which do not pass through any other point in W. If a line segment links two points in U, its slope must be rational or ∞, hence it would not

pass through any point in V. Furthermore, it does not pass through any other point in U if and only if its holonomy is a pair of coprime integers. The same is true for line segments linking two points in V. On the other hand, given any point $p \in U$ and any point $q \in V$, a line segment from p to q has irrational slope hence cannot pass through any other point in U or V, therefore the holonomy of such line segment can be any vector in $\mathbb{Z}^2+(\sqrt{2}-1, \sqrt{3}-1)$. Similarly the holonomies of saddle connections from V to U are $\mathbb{Z}^2 - (\sqrt{2}-1, \sqrt{3}-1)$. Hence $S_{M_1} = \{(a,b) \in \mathbb{Z}^2 : \gcd(a,b) = 1\} \cup (\mathbb{Z}^2+(\sqrt{2}-1, \sqrt{3}-1)) \cup (\mathbb{Z}^2-(\sqrt{2}-1, \sqrt{3}-1))$, this set is uniformly discrete, and the last two pieces are uniformly dense.

Questions:

(1) Can the set of holonomy vectors of all saddle connections of a non-arithmetic lattice surface be relatively dense?
(2) Is there any characterization of the set of flat surfaces M such that S_M are Deloné, relatively dense or uniformly discrete?
(3) Is there a surface M which is not a branched cover of the torus for which S_M is Deloné?

References

[AC12] Jayadev S Athreya and Jon Chaika. The distribution of gaps for saddle connection directions. *Geometric and Functional Analysis*, 22(6):1491–1516, 2012.

[ACL13] Jayadev S Athreya, Jon Chaika, and Samuel Lelievre. The gap distribution of slopes on the golden l. *arXiv preprint arXiv:1308.4203*, 2013.

[EM01] Alex Eskin and Howard Masur. Asymptotic formulas on flat surfaces. *Ergodic Theory and Dynamical Systems*, 21(02):443–478, 2001.

[HS71] Fritz Herzog and BM Stewart. Patterns of visible and nonvisible lattice points. *American Mathematical Monthly*, pages 487–496, 1971.

[KS00] Richard Kenyon and John Smillie. Billiards on rational-angled triangles. *Commentarii Mathematici Helvetici*, 75(1):65–108, 2000.

[Mas88] Howard Masur. Lower bounds for the number of saddle connections and closed trajectories of a quadratic differential. In *Holomorphic Functions and Moduli I*, pages 215–228. Springer, 1988.

[Mas90] Howard Masur. The growth rate of trajectories of a quadratic differential. *Ergodic Theory and Dynamical Systems*, 10(01):151–176, 1990.

[Sen06] Marjorie Senechal. What is a quasicrystal. *Notice of the AMS*, 53:886–887, 2006.

[SW10] John Smillie and Barak Weiss. Characterizations of lattice surfaces. *Inventiones mathematicae*, 180(3):535–557, 2010.

[Vee89] William A Veech. Teichmüller curves in moduli space, Eisenstein series and an application to triangular billiards. *Inventiones mathematicae*, 97(3):553–583, 1989.